

## **Hydrological Sciences Journal Contents 2007**

### **Papers published in 2007**

- Editorial—Quantifying the impact of hydrological studies *Demetris Koutsoyiannis & Zbigniew W. Kundzewicz* **52(1)**, 3–17.
- Synthèse des techniques d'homogénéisation des séries climatiques et analyse d'applicabilité aux séries de précipitations *Claudie Beaulieu, Taha B. M. J. Ouarda & Ousmane Seidou* **52(1)**, 18–37.
- Modelling the impact of climatic variability on water resources in West and Central Africa from a non-calibrated hydrological model *J. E. Paturel, C. Barrau, G. Mahé, A. Dezetter & E. Servat* **52(1)**, 38–48.
- Estimation des pluies exceptionnelles journalières en zone tropicale: cas de la Côte d'Ivoire par comparaison des lois lognormale et de Gumbel *Albert Goula Bi Tie, Brou Konan, Yao Téléphore Brou, Savané Issiaka, Vamoryba Fadika & Bernard Srohourou* **52(1)**, 49–67.
- Un siècle de débits annuels du fleuve Sénégal *Pierre Hubert, Jean-Claude Bader & Hocine Bendjoudi* **52(1)**, 68–73.
- Long-term changes in river temperature and the influence of climatic and hydrological factors *Bruce W. Webb & Franz Nobilis* **52(1)**, 74–85.
- Estimation of an index flood using data transfer in the UK *Thomas Rodding Kjeldsen & David Jones* **52(1)**, 86–98.
- Monthly dam inflow forecasts using weather forecasting information and neuro-fuzzy technique *Deg-Hyo Bae, Dae Myung Jeong & Gwangseob Kim* **52(1)**, 99–113.
- Multi-step-ahead neural networks for flood forecasting *Fi-John Chang, Yen-Ming Chiang & Li-Chiu Chang* **52(1)**, 114–130.
- Impact of limited streamflow data on the efficiency and the parameters of rainfall–runoff models *Charles Perrin, Ludovic Oudin, Vazken Andréassian, Claudia Rojas-Serna, Claude Michel & Thibault Mathevet* **52(1)**, 131–151.
- A study of dynamical seasonal prediction of potential water resources based on an atmospheric GCM experiment with prescribed sea-surface temperature *Tosiyuki Nakaegawa, Shoji Kusunoki, Masato Sugi, Akio Kitoh, Chiaki Kobayashi & Kiyoharu Takano* **52(1)**, 152–165.
- Long-term freshwater inflow and sediment discharge into Lake Pontchartrain in Louisiana, USA *Kangsheng Wu & Y. Jun Xu* **52(1)**, 166–180.
- Sedimentation rate and pattern of a Himalayan foothill lake using  $^{137}\text{Cs}$  and  $^{210}\text{Pb}$  *S. P. Rai, Vijay Kumar & Bhishm Kumar* **52(1)**, 181–191.
- Hydrodynamic behaviour of karstic aquifers in Boroujerd, western Iran *Babak Ebrahimi, Mehrdad Pasandi & Mohammad R. Ahmadipour* **52(1)**, 192–205.
- Evidence of karst from behaviour of the Asmari limestone aquifer at the Khersan 3 Dam site, southern Iran *Zargham Mohammadi, Ezzatollah Raeisi & Michel Bakalowicz* **52(1)**, 206–220.
- Steady-state groundwater flow model with variable hydraulic conductivity *Abdüsselam Altunkaynak* **52(1)**, 221–229.
- On width function-based unit hydrographs deduced from separately random self-similar river networks and rainfall variability. Discussion of “Coding random self-similar river networks and calculating geometric distances: 1. General methodology and 2. Application to runoff simulations” by C.-P. Hung & R.-Y. Wang, *Hydrol. Sci. J.* **50(5)**, 753–768 and 769–781. *Christophe Cudennec* **52(1)**, 230–237.
- A dynamic random self-similar river network during the rainfall–runoff process. Reply to Discussion by C. Cudennec. *Chun-Po Hung & Ru-Yih Wang* **52(1)**, 237–244.
- Future long-term changes in global water resources driven by socio-economic and climatic changes *Joseph Alcamo, Martina Flörke & Michael Märker* **52(2)**, 247–275.
- Long-term variations in orographic rainfall: analysis and implications for upland catchments *A. R. Malby, J. D. Whyatt, R. J. Timmis, R. L. Wilby & H. G. Orr* **52(2)**, 276–291.

- Predicting daily streamflow in ungauged rural catchments: the case of Masinga catchment, Kenya *Benedict Mwavu Mutua & Andreas Klik* **52**(2), 292–304.
- Interpolating monthly precipitation by self-organizing map (SOM) and multilayer perceptron (MLP) *Aman Mohammad Kalthé & Ronny Berndtsson* **52**(2), 305–317.
- Intégration d'un filtre de Kalman dans le modèle hydrologique HBV pour la prévision des débits *Rim Ouachani, Zoubeida Bargaoui & Taha Ouarda* **52**(2), 318–337.
- A new drain spacing formula *G. C. Mishra & Vivekanand Singh* **52**(2), 338–351.
- Optimal pumping locations of skimming wells *S. V. N. Rao & S. Manju* **52**(2), 352–361.
- Critical appraisal of BOD and DO models applied to a highly polluted river in India *Ramakar Jha, C. S. P. Ojha & K. K. S. Bhatia* **52**(2), 362–375.
- Snow and glacier melt contribution in the Beas River at Pandoh Dam, Himachal Pradesh, India *Vijay Kumar, Pratap Singh & Vivekanand Singh* **52**(2), 376–388.
- Hydroinformatics: computational intelligence and technological developments in water science applications—Editorial *Linda See, Dimitri Solomatine, Robert Abraham & Elena Toth* **52**(3), 391–396.
- Multi-objective performance comparison of an artificial neural network and a conceptual rainfall–runoff model *N. J. de Vos & T. H. M. Rientjes* **52**(3), 397–413.
- Timing error correction procedure applied to neural network rainfall–runoff modelling *Robert J. Abraham, Alison J. Heppenstall & Linda M. See* **52**(3), 414–431.
- Comparative assessment of six automatic optimization techniques for calibration of a conceptual rainfall–runoff model *Monomoy Goswami & Keiran M. O'Connor* **52**(3), 432–449.
- Watershed calibration using multistart local optimization and evolutionary optimization with radial basis function approximation *Christine A. Shoemaker, Rommel G. Regis & Ryan C. Fleming* **52**(3), 450–465.
- Development of a low-flow forecasting model using the M5 machine learning method *Luka Stravs & Mitja Brilly* **52**(3), 466–477.
- Fuzzy logic-based rainfall–runoff modelling using soil moisture measurements to represent system state *Markus Casper, Peter Gemmar, Oliver Gronz, Margret Johst & Manfred Stüber* **52**(3), 478–490.
- Baseflow separation techniques for modular artificial neural network modelling in flow forecasting *Gerald Corzo & Dimitri Solomatine* **52**(3), 491–507.
- Seasonal reservoir inflow forecasting with low-frequency climatic indices: a comparison of data-driven methods *Getnet Muluye & Paulin Coulibaly* **52**(3), 508–522.
- A probabilistic model to support reservoir operation decisions during flash floods *L. Mediero, L. Garrote & F. Martín-Carrasco* **52**(3), 523–537.
- Simulation of the hydrological processes on reconstructed watersheds using system dynamics *Amin Elshorbagy, A. Jutla & J. Kells* **52**(3), 538–562.
- Modelling the dynamics of the evapotranspiration process using genetic programming *Kamban Parasuraman, Amin Elshorbagy & Sean K. Carey* **52**(3), 563–578.
- Cellular automata algorithms for drainage network extraction and rainfall data assimilation *Erika Coppola, Barbara Tomassetti, Laura Mariotti, Marco Verdecchia & Guido Visconti* **52**(3), 579–592.
- Analysis and propagation of uncertainties due to the stage–discharge relationship: a fuzzy set approach *Rajesh Raj Shrestha, Andrés Bárdossy & Franz Nestmann* **52**(4), 595–610.
- To prewhiten or not to prewhiten in trend analysis? *M. Bayazit & B. Önöz* **52**(4), 611–624.
- A national low flow estimation procedure for Austria *Gregor Laaha & Günter Blöschl* **52**(4), 625–644.
- The spatially heterogeneous distribution of precipitation in the Anduo area, Tibetan Plateau, in summer 1998 *Meixue Yang, Tandong Yao, Xiaohua Gou, Huijun Wang & Hongguan Tang* **52**(4), 645–653.
- Modélisation du bilan hydrique du bassin versant du Lac Ba Bê, Viêt-nam, à l'aide de données physiographiques issues de télédétection et de données climatologiques *Laurie St Onge, Ferdinand Bonn, Alain N. Rousseau, Pham Van Cu & Nguyen Quang My* **52**(4), 654–670.
- The characteristics of flood responses to the restoration of polders on Dongting Lake, China *Jiang Jiahu, Lai Xijun & Huang Qun* **52**(4), 671–685.
- Optimal reservoir operation for irrigation of multiple crops using elitist-mutated particle swarm optimization *M. Janga Reddy & D. Nagesh Kumar* **52**(4), 686–701.
- The *sudestadas*: a hydro-meteorological phenomenon that affects river pollution (River Luján, South America) *Haydée Pizarro, Patricia Rodríguez, Stella Maris Bonaventura, Inés O'Farrell & Irina Izaguirre* **52**(4), 702–712.

- Comparing artificial neural networks and regression models for predicting faecal coliform concentrations *Diane M. L. Mas & David P. Ahlfeld* **52**(4), 713–731.
- Investigation of a chalk aquifer: advantages and limits of diffusion samplers and low-flow sampling techniques *Frédéric Barrez, Jacky Mania, Jean-Loius Mansy & Francis Robaszynski* **52**(4), 732–747.
- Quantification of heterogeneity of the unsaturated zone based on environmental deuterium observed in lysimeter experiments *Christine Stumpp, Piotr Maloszewski & Willibald Stichler* **52**(4), 748–762.
- Explicit algebraic influence coefficients: a one-dimensional transient aquifer model *Mohamed Saffi & Abdelkhalek Cheddadi* **52**(4), 763–776.
- Trends in suspended sediment grain size in the upper Yangtze River and its tributaries, as influenced by human activities *Jiongxin Xu* **52**(4), 777–792.
- Deriving stage–discharge–sediment concentration relationships using fuzzy logic *A. K. Lohani, N. K. Goel & K. K. S. Bhatia* **52**(4), 793–807.
- Hyperspectral feature analysis of chlorophyll *a* and suspended solids with field measurements in Taihu Lake, eastern China *Ronghua Ma, Xiaodong Ma & Jinfang Dai* **52**(4), 808–824.
- Discussion of “Generalized regression neural networks for evapotranspiration modelling” by O. Kişi, *Hydrol. Sci. J.* **51**(6), 1092–1105. *Hafzullah Aksoy, Aytac Guven, Ali Aytekin, M. Ishak Yucesu & N. Erdem Unal* **52**(4), 825–828.
- Reply to Discussion by Aksoy *et al.* *Ozgür Kişi* **52**(4), 829–831.
- Discussion of “Generalized regression neural networks for evapotranspiration modelling” by O. Kişi, *Hydrol. Sci. J.* **51**(6), 1092–1105. *Demetris Koutsoyiannis* **52**(4), 832–835.
- Reply to Discussion by D. Koutsoyiannis. *Ozgür Kişi* **52**(4), 836–839.
- Rainfall–runoff response, event-based runoff coefficients and hydrograph separation *Theresa Blume, Erwin Zehe & Axel Bronstert* **52**(5), 843–862.
- Probability distribution of annual, seasonal and monthly precipitation in Japan *Sheng Yue & Michio Hashino* **52**(5), 863–877.
- The MGB-IPH model for large-scale rainfall–runoff modeling *Walter Collischonn, Daniel Allasia, Benedito C. Da Silva & Carlos E. M. Tucci* **52**(5), 878–895.
- A comparative study of three neural network forecast combination methods for simulated river flows of different rainfall–runoff models *Asaad Y. Shamseldin, Kieran M. O’Connor & Ahmed E. Nasr* **52**(5), 896–916.
- Simulation and spatio-temporal disaggregation of multi-site rainfall data for urban drainage applications *Marie-Laure Segond, Natasa Neokleous, Christos Makropoulos, Christian Onof & Cedo Maksimovic* **52**(5), 917–935.
- Measured effects of new lake surfaces on regional precipitation *Tobias Conradt, Zbigniew W. Kundzewicz, Fred Hattermann & Frank Wechsung* **52**(5), 936–955.
- Regional tests for trend detection in maximum precipitation series in the French Mediterranean region *Nicolas Pujol, Luc Neppel & Robert Sabatier* **52**(5), 956–973.
- Non-identical models for seasonal flood frequency analysis *Bin Fang, Shenglian Guo, Shanxu Wang, Pan Liu & Yi Xiao* **52**(5), 974–991.
- Assessment of severity of the extreme River Tyne flood in January 2005 using gauged and historical information *David R. Archer, Francois Leesch & Kirsty Harwood* **52**(5), 992–1003.
- Flooding in Fiji: findings from a 100-year historical series *Stephen W. Yeo, Russell J. Blong & K. John McAneney* **52**(5), 1004–1015.
- Coping with floods: preparedness, response and recovery of flood-affected residents in Germany in 2002 *Annegret H. Thieken, Heidi Kreibich, Meike Müller & Bruno Merz* **52**(5), 1016–1037.
- Optimal contour mapping of groundwater levels using universal kriging—a case study *Vijay Kumar* **52**(5), 1038–1050.
- Groundwater recharge estimation using improved soil moisture balance methodology for a tropical climate with distinct dry seasons *C. Shanthy de Silva & Ken R. Rushton* **52**(5), 1051–1067.
- Applicability of the Modified Universal Soil Loss Equation for prediction of sediment yield in Khanmirza watershed, Iran *S. H. R. Sadeghi & T. Mizuyama* **52**(5), 1068–1075.
- Dryland hydrology in Mediterranean regions—a review *Christophe Cudennec, Christian Leduc & Demetris Koutsoyiannis* **52**(6), 1077–1087.
- Structure du gradient pluviométrique de la transition Méditerranée–Sahara en Tunisie: déterminants géographiques et saisonnalité *Mohamed Slimani, Christophe Cudennec & Haïfa Feki* **52**(6), 1088–1102.

- Regression analysis of rainfall–runoff data from an arid catchment in Oman *Neil McIntyre, Aisha Al-Qurashi & Howard Wheeler* **52**(6), 1103–1118.
- Hydrological impact simulations of climate change on Lebanese coastal rivers *Antoine Hreiche, Wajdi Najem & Claude Bocquillon* **52**(6), 1119–1133.
- Caractéristiques et impacts hydrologiques de banquettes en cascade sur un versant semi-aride en Tunisie centrale *Slah Nasri* **52**(6), 1134–1145.
- Hydrological implications of desertification in southeastern Spain *Wenceslao Martín-Rosales, Antonio Pulido-Bosch, Ángela Vallejos, Juan Gisbert, José Miguel Andreu, & Francisco Sánchez-Martos, F.* **52**(6), 1146–1161.
- Impacts of hydrological changes in the Mediterranean zone: environmental modifications and rural development in the Merguellil catchment, central Tunisia *C. Leduc, S. Ben Ammar, G. Favreau, R. Béji, R. Virrion, G. Lacombe, J. Tarhouni, C. Aouadi, B. Zenati Chelli, N. Jebnoun, M. Oï, J.-L. Michelot, & K. Zouari* **52**(6), 1162–1178.
- La gestion communautaire de l'eau agricole à l'épreuve des politiques participatives: regards croisés Tunisie/Maroc *Bruno Romagny & Jeanne Riaux* **52**(6), 1179–1198.
- Regionalizing fine time-scale rainfall affected by topography in semi-arid Tunisia *Sihem Jebari, Ronny Berndtsson, Cintia Uvo & Akissa Bahri* **52**(6), 1199–1215.
- Hydrological impacts of changes in evapotranspiration and precipitation: two case studies in semi-arid and humid climates *Markus Möller & Gerald Stanhill* **52**(6), 1216–1231.
- Modélisation hydrologique stochastique d'un bassin affecté par des changements d'occupation: cas du Merguellil en Tunisie centrale *Ahmadi Kingumbi, Zoubeida Bargaoui, Emmanuel Ledoux, Mustapha Besbes & Pierre Hubert* **52**(6), 1232–1252.
- Modelling the effects of a bushfire on erosion in a Mediterranean basin *Gian Vito di Piazza, Costanza di Stefano & Vito Ferro* **52**(6), 1253–1270.
- Key processes influencing erosion and sediment transport in a semi-arid Mediterranean area: the upper Tafna catchment, Algeria *Abdesselam Megnounif, Abdelali Terfous, Abdellah Ghenaim & Jean-Bernard Poulet* **52**(6), 1271–1284.
- Elaboration d'un modèle d'érosion hydrique à la parcelle (PLAG) et son application pour l'estimation des apports en sédiments en zone semi-aride tunisienne *Lamia Lajili-Ghezal* **52**(6), 1285–1297.
- Etude intégrée des données de surface et de subsurface pour la prospection des bassins hydrogéothermiques: cas du bassin de Maknassy (Tunisie centrale) *S. Bourri, M. Gasmî, M. Jaouadi, I. Souissi, A. Lahlou Mimi & H. Ben Dhia* **52**(6), 1298–1315.
- Lake Victoria: derivation of a corrected natural water level series *J. V. Sutcliffe & G. Petersen* **52**(6), 1316–1321.

### Books reviewed in 2007

- Les eaux continentales* sous la direction de G. de Marsily **52**(1), 245–246.
- Living with Nature's Extremes – The Life of Gilbert Fowler White* by Robert E. Hinshaw **52**(4), 840–841.
- Flood Risk Management in Europe – Innovation in Policy and Practice* edited by Selina Begum, Marcel Stive & Jim Hall **52**(4), 841–842.

## Key words index 2007

- abrupt change, 68
- Africa, 38
- Algeria, 1271
- alpha-level cut, 595
- anthropogenic change, 1162
- aquifer system, 206
- aquifers, 1232
- ArcView GIS, 292
- arid zone, 1103
- artificial neural network(s), 352, 397, 563, 713, 793
  - forecasting, 491
- assessment, 1285
- Atlantic mountain range, 1119
- Austria, 74
  
- baseflow, 192, 843
  - separation, 491
- Bayesian
  - methods, 18
  - networks, 523
  - neural networks, 508
- biochemical oxygen demand (BOD), 362
  
- caesium-137, 181
- calibration, 38, 131
- catchment grouping, 625
- cellular automata, 579
- central Tunisia, 1134, 1232, 1298
- chalk aquifer, 732
- Chile, 843
- China, 671
- chlorophyll, 808
- Churchill Falls basin, 508
- CHYM, 579
- class A pan, 1216
- climate
  - and hydrology, 74
  - change, 276, 917, 1119
    - and water, 247
  - variability adjustment, 625
- climatic
  - change, 956
  - scenario, 38
  - series, 18
  - variability, 38
- coliform concentration, 713
- compartment, 763
- conceptual models, 397
- contour ridges, 1134
- correlation, 362
- coupled
  - hydrodynamic model, 671
  - surface water-groundwater flow model, 1232
- crop
  - water allocation, 686
  - yield response, 686
- cumulative rainfall, 376
  
- daily
  - rainfall, 1119
  - streamflow, 292
- dam inflow, 99
- Darcy law, 221
- data
  - assimilation, 318
  - transfer, 86
  - driven
    - methods, 508
    - models, 491
- deoxygenation, 362
- desertification, 1146
- dialysis membrane, 732
- disaggregation, 917
- discharge, 68, 793
- dissolved oxygen (DO), 362
- Dongting Lake, 671
- drainage network extraction, 579
- drought(s), 68, 625
- dynamical seasonal prediction, 152
  
- eddy-covariance, 563
- efficiency of entrainment, 1285
- elitist mutation, 686
- ENSO, 508
- environmental isotopes, 206, 748
- equivalent depth, 338
- erosion, 1134
- evaporation, 152
- evapotranspiration, 563, 1051
  - losses, 376
- event-based runoff coefficient, 843
- evolutionary algorithm, 450
- extension principle, 595
- extreme
  - flood, 992
  - rainfall, 49
  
- Fiji, 1004
- fine time-scale rainfall, 1119
- fire, 1253
- flood
  - forecasting, 114
  - frequency, 1004
  - impact, 1016
  - response, 671
  - risk assessment, 992
  - warning, 1016
- flooding, 936

- frequency analysis, 974  
 freshwater, 166  
 fuzzy  
   inference system, 793  
   logic, 478  
   number, 595  
   regression analysis, 595
- GAME-Tibet, 645  
 generalized  
   value distribution, 956  
   linear models, 917  
 genetic  
   algorithm, 432  
   programming, 563  
 geochemistry, 1162  
 geomorphology-based unit hydrograph, 230  
 geostatistics, 1088  
   semivariogram, 1038  
 geothermal gradient, 1298  
 geothermometry, 1298  
 Germany, 1016  
 glacier melt, 376  
 global  
   change, 247  
   climate model, 152  
   optimization, 450  
   water resources, 247  
 gravel extraction, 992  
 Green function, 763  
 groundwater, 221, 352  
   levels, 1038  
   sampling, 732  
 Gulf of Mexico, 166  
 Gumbel distribution, 49
- HBV model, 318  
 heterogeneity, 748  
 historical data, 992  
 Holdridge classification, 38  
 homogenization, 18  
 Hooghoudt's drain spacing formula, 338  
 Hubert segmentation, 68  
 humid tropical zone, 49  
 hydrochemical stratification, 732  
 hydrochemistry, 206  
 hydrodynamic model, 595  
 hydrodynamics, 1162  
 hydrogeothermal system, 1298  
 hydrograph separation, 491, 843  
 hydrological  
   forecasting, 414  
   model, 579, 878  
   modelling, 38, 292  
 hyperspectral reflectance, 808  
 hysteresis, 1271
- identification, 763  
 illegal pumping, 763  
 index flood, 86  
 India, 181  
 influence coefficient, 763  
 institutional dynamics, 1179
- inter-annual variability, 1216  
 interpolation, 305, 917  
 Iran, 192, 206, 1068  
 irrigated systems, 1179  
 irrigation  
   of multiple crops, 686  
   water requirement, 1216  
 Ivory Coast, 49
- Jammu and Kashmir, 181  
 Japan, 863  
 jump, 18
- Kalman filter, 318  
 karst, 192, 206  
 Khanmirza watershed, 1068  
 Khersan 3 Dam, 206
- Lake  
   District, England, 276  
   Pontchartrain, 166  
   restoration, 671  
   Victoria, 1316  
 Lamb weather types, 276  
 land-use change, 936, 1232  
 large basins, 878  
 lead-210, 181  
 linear  
   statistical model, 843  
   trends, 1216  
 L-moment, 863  
 local optimization, 450  
 lognormal distribution, 49  
 long-term trends, 74  
 low flows, 466, 625  
   forecasting model, 466  
   sampling, 732  
 lumped parameter model, 748  
 Lusatia, 936
- M5, 466  
 machine learning, 466  
 Maghreb, 1179  
 Mann-Kendall test, 611, 956  
 Mansar Lake, 181  
 mapping, 49  
 Masinga catchment, Kenya, 292  
 maximum-likelihood method, 956  
 mean transit time, 748  
 mechanisms of erosion, 1271  
 Mediterranean, 1119  
   basin, 1253  
 missing data, 305  
 mitigation, 1016  
 model structure, 230  
 modeling, 563, 1103, 1253, 1285  
 modular models, 491  
 multilayer perceptron (MLP), 305, 896  
 multi-level single linkage, 450  
 multi-objective calibration, 397  
 multistart, 450  
 multi-step-ahead, 114  
 multivariate analysis, 1119

- MUSLE, 1068
- naturalised lake levels, 1316
- neural network, 114, 414  
 combination methods, 896
- neuro-fuzzy system, 99
- non-identical model, 974
- nonlinear process, 793
- non-point source pollution, 362
- nonstationarity, 68
- North Atlantic Oscillation, 74
- northern  
 Iran, 305  
 Qinghai-Xizang (Tibetan) Plateau, 645
- optimization, 432
- overexploitation, 1146
- parameter fitting, 878
- participative management, 1179
- particle swarm optimization, 432, 686
- Penman-Monteith method, 563
- plane, 1285
- Poisson-cluster processes, 917
- pooled average relative error, 793
- population growth, 166
- POT sampling, 974
- potential predictability, 152
- power of a test, 611
- precipitation, 152, 305, 863, 936
- prediction of ungauged catchments (PUB), 625
- predictive uncertainty, 625
- preferential flow, 748
- prewhitening, 611
- principal component analysis, 936
- probabilistic forecast, 523
- probability distribution, 863
- public policies, 1179
- quickflow, 192
- radial basis function, 450, 896
- rainfall, 917  
 -generating trajectory, 1088  
 gradient, 1088  
 space-time variability, 230  
 917
- rainfall-runoff, 1103  
 model, 230, 896  
 modelling, 131, 318, 414, 478, 491  
 response, 843
- rainshadow, 276
- re-aeration, 362
- real time, 318
- recession  
 analysis, 466  
 curve, 192
- recharge, 1051
- reconstructed watersheds, 538
- recovery, 1016
- recurrent networks, 508
- regional  
 climatic change, 936  
 regression, 625  
 test for trend detection, 956
- regionalization, 625, 1119
- regression, 713, 793, 1103  
 trees, 466
- relief, 1088
- remote sensing, 808
- reservoir, 777  
 inflows, 508  
 operation, 523, 686
- return period, 1216
- Río de la Plata Estuary, 702
- River  
 flow simulation, 896  
 Luján, 702  
 network, 230  
 Ouse, 414  
 Taquari, 878  
 temperature, 74  
 Tyne (UK), 992  
 Uruguay, 878
- Rosenbrock, 432
- runoff, 1051, 1134  
 forecasting, 397
- sampling, 131
- seasonal  
 flood, 974  
 forecast, 508  
 runoff, 152
- seasonality, 1088  
 index, 625
- sediment  
 grain size, 777  
 graph, 1068  
 rating curve, 793  
 yield, 1068, 1253
- sedimentation rate, 181
- sedimentogram, 1285
- self-organizing map (SOM), 305
- semi-arid, 1119  
 area, 1271  
 region, 1134
- semi-confined aquifer, 763
- Senegal, 68
- sensitivity analysis, 131
- serial  
 correlation, 611  
 -propagated structure, 114
- shuffled complex evolution, 432
- simple neural network, 896
- simplex, 432
- simulated annealing, 352, 432
- simulations, 1119
- singularity, 68
- skimming wells, 352
- SMAR, 432
- snow, 1119  
 covered area, 376
- snowmelt, 376
- SOI, 1004
- soil  
 conservation, 777

- erodibility, 1253
- hydrophobicity, 1253
- moisture, 478
  - balance, 1051
  - dynamics, 686
  - properties, 1051
- South America, 702, 878
- Spatial
  - heterogeneous distribution, 645
  - rainfall patterns, 1119
- spring, 192
- Sri Lanka, 1051
- stage, 793
- stage–discharge relationship, 595
- statistical
  - hydrology, 49
  - test, 18
- steady-state flow, 221
- Stream Flow Model, 292
- streamflow
  - data, 131
  - impact, 1119
- Subsurface drain spacing, 338
- subtractive clustering, 99
- sudestadas, 702
- summer monsoon precipitation, 645
- suspended
  - sediment load, 777, 1271
  - solids, 808, 166
- SWAT, 450
- system dynamics, 538
  
- Taihu Lake, China, 808
- Taiwan, 114
- temporal trends, 1004
- thermal spring, 1298
- Thornthwaite & Mather method, 654
- thunderstorm activity, 936
- time domain reflectometry (TDR), 478
- time series analysis, 68
- time-lagged networks, 508
- timing errors, 414
- trend(s), 18, 1146
  - analysis, 611
- tropical cyclones, 1004
- Tunisia, 1119, 1088, 1162
  - central, 1134, 1232, 1298
- Tunisian semi-arid region, 1285
- type curve, 221
  
- uncertainty, 86, 595
  - ungauged
    - basins, 878
    - catchment, 292
    - site, 86
- universal kriging, 1038
- unsaturated zone, 748
- unsteady state, 763
- upcoming, 352
- updating, 318
- upper Yangtze River, 777
  
- variable hydraulic conductivity, 221
  
- Vietnam, 654
- vulnerability, 748
  
- Wadi, 1103
  - Sebdou, 1271
- Water
  - availability, 247
  - balance, 654, 936, 1162
    - approach, 376
    - model, 1216
  - erosion, 1285
  - pollution, 702
  - quality, 713
  - resources, 152, 1232
  - scenarios, 247
  - shortage, 1146
  - stress, 247
  - user associations, 1179
- watershed, 654, 1232
  - calibration, 450
  - modelling, 538
- weather forecasting information, 99
  
- West Africa, 68
- width function, 230
  
- Zagros, 206

## Index des mots clefs 2007

- activité orageuse, 936  
 afflux dans un barrage, 99  
 Afrique, 38  
   de l'Ouest, 68  
 ajustement  
   de variabilité climatique, 625  
   des paramètres, 878  
 alerte de crue, 1016  
 Algérie, 1271  
 Algorithme  
   évolutif, 450  
   génétique, 432  
 Allemagne, 1016  
 allocation de l'eau agricole, 686  
 alpha-coupe, 595  
 Amérique du Sud, 702, 878  
 Analyse  
   de récession, 466  
   de régression floue, 595  
   de sensibilité, 131  
   de séries temporelles, 68  
   de tendance, 611  
   en composantes principales, 936  
   fréquentielle, 974  
   multivariable, 1119  
 apprentissage machine, 466  
 approche du bilan hydrologique, 376  
 aquifère(s), 1232  
   de craie, 732  
   semi-confiné, 763  
 arbres de régression, 466  
 assimilation des données, 318  
 associations d'usagers de l'eau, 1179  
 automates cellulaires, 579  
 Autriche, 74  
  
 bac de classe A, 1216  
 banquettes, 1134  
 barrage, 777  
   de Khersan 3, 206  
  
 bassin(s) versant(s), 1232  
   de Churchill Falls, 508  
   de Khanmirza, 1068  
   de Masinga, Kenya, 292  
   non jaugé(s), 292, 878  
   reconstruits, 538  
   grands, 878  
 besoin en eau d'irrigation, 1216  
 bilan  
   d'humidité du sol, 1051  
   hydrique, 654, 936  
   hydrologique, 1162  
  
 calage, 38, 131  
   au niveau du bassin versant, 450  
 calibration multi-objective, 397  
  
 carte auto-organisée (SOM), 305  
 cartographie, 49  
 césium-137, 181  
 chaîne atlasique, 1119  
 changement  
   anthropique, 1162  
   climatique  
     et eau, 247  
     régional, 936  
     changement climatique, 276,  
     917, 956, 1119  
   d'occupation du sol, 936, 1232  
   global, 247  
 charge de sédiments en suspension, 777, 1271  
 Chili, 843  
 Chine, 671  
 chlorophylle, 808  
 CHYM, 579  
 Classification  
   de Holdridge, 38  
   soustractive, 99  
 climat et hydrologie, 74  
 coefficient  
   d'écoulement événementiel, 843  
   d'influence, 763  
 compartiment, 763  
 concentration en coliformes, 713  
 conductivité hydraulique variable, 221  
 conservation des sols, 777  
 corrélation, 362  
   sérielle, 611  
 Côte d'Ivoire, 49  
 courbe  
   de décrue, 192  
   de production sédimentaire, 793  
   type, 221  
 covariances turbulentes, 563  
 croissance démographique, 166  
 crue  
   extrême, 992  
   saisonniers, 974  
 cyclones tropicaux, 1004  
  
 débit, 68, 793  
   de base, 843  
 déficit hydrique, 1146  
 demande biochimique en oxygène (DBO), 362  
 démarrage multiple, 450  
 désagrégation, 917  
 désertification, 1146  
 désoxygénation, 362  
 disponibilité de l'eau, 247  
 distribution  
   de probabilité, 863  
   généralisée des valeurs extrêmes, 956  
   spatiale hétérogène, 645

- données
  - de débit, 131
  - historiques, 992
  - manquantes, 305
- dynamique
  - de l'humidité du sol, 686
  - des systèmes, 538
  - institutionnelle, 1179
- eau
  - douce, 166
  - souterraine, 352
- échantillonnage, 131
  - bas débit, 732
  - de l'eau souterraine, 732
  - POT, 974
- écoulement(s)
  - de base, 192
  - journalier, 292
  - préférentiel, 748
  - quasi-stationnaire, 221
  - rapide, 192
  - saisonnier, 152
  - entrant dans un réservoir, 508
- efficacité d'entraînement, 1285
- ENSO, 508
- érodibilité des sols, 1253
- érosion, 1134
  - hydrique, 1285
- erreur(s)
  - relative moyenne d'ensemble, 793
  - temporelles, 414
- espacement de drains de subsurface, 338
- estimation, 1285
- Estuaire du Río de la Plata, 702
- étiages, 625
- évaluation du risque d'inondation, 992
- évaporation, 152
- évapotranspiration, 563, 1051
- extraction
  - de gravier, 992
  - du réseau de drainage, 579
- faibles débits, 466
- Fidji, 1004
- filtre de Kalman, 318
- Fleuve Yangtze supérieur, 777
- fonction(s)
  - de base radiale, 450, 896
  - de Green, 763
  - largeur, 230
- fonte
  - glaciaire, 376
  - nivale, 376
- formule d'espacement de drains de Hooghoudt, 338
- fréquence des crues, 1004
- GAME-Tibet, 645
- géochimie, 1162
- géostatistique, 1038, 1088
- géothermométrie, 1298
- gestion
  - de réservoir, 686
  - participative, 1179
- Golfe de Mexico, 166
- gradient
  - géothermique, 1298
  - pluviométrique, 1088
- grands bassins versants, 878
- granulométrie sédimentaire, 777
- hauteur, 793
- hétérogénéité, 748
- homogénéisation, 18
- humidité du sol, 478
- hydrodynamique, 1162
- hydrogramme
  - unitaire à base géomorphologique, 230
  - séparation événementielle, 843
- hydrologie statistique, 49
- hydrophobicité des sols, 1253
- hystérésis, 1271
- identification, 763
- impact
  - des crues, 1016
  - sur les écoulements, 1119
- incendie, 1253
- incertitude, 86, 595
  - de prévision, 625
- Inde, 181
- indice
  - de crue, 86
  - de saisonnalité, 625
- informations de prévision météorologique, 99
- inondation, 936
- interpolation, 305, 917
- Iran, 192, 206, 1068
- irrigation de cultures multiples, 686
- isotopes
  - du milieu, 206
  - environnementaux, 748
- Jammu et Kashmir, 181
- Japon, 863
- karst, 206
- krigeage universel, 1038
- Lac
  - Dongting, 671
  - Mansar, 181
  - Ponchartrain, 166
  - Taihu, Chine, 808
  - Victoria, 1316
- Lake District, Angleterre, 276
- L-moment, 863
- logique floue, 478
- loi
  - de Darcy, 221
  - de Gumbel, 49
  - lognormale, 49
- Lusatia, 936

- M5, 466  
 Maghreb, 1179  
 Matières  
   en suspension, 166  
   solides en suspension, 808  
 mécanismes d'érosion, 1271  
 Méditerranée, 1119  
 membrane de dialyse, 732  
 méthode(s)  
   de Pennman-Monteith, 563  
   de Thornthwaite & Mather, 654  
   Bayésiennes, 18  
   conditionnées par les données, 508  
   de combinaison de réseaux de neurones, 896  
 milieu semi-aride, 1134  
 mise à jour, 318  
 modelage, 1253  
 modèle(s)  
   climatique global, 152  
   conceptuel, 397  
   conditionnés par les données, 491  
   couplé eaux de surface-eaux souterraines, 1232  
   de bilan hydrologique, 1216  
   de prévision des faibles débits, 466  
   global, 748  
   HBV, 318  
   hydrodynamique, 595  
     couplé, 671  
   hydrologique, 579, 878  
   linéaires généralisés, 917  
   modulaires, 491  
   non-identique, 974  
   pluie-débit, 896  
   statistique linéaire, 843  
 modélisation, 563, 1285, 1103  
   de bassin versant, 538  
   hydrologique, 38, 292  
   pluie-débit, 131, 230, 318, 414, 478, 491  
 multi-échéances, 114  
 MUSLE, 1068  
 mutation élitiste, 686  
  
 neige, 1119  
 niveaux piézométriques, 1038  
 nombre flou, 595  
 non stationnarité, 68  
 nord  
   de l'Iran, 305  
   du Plateau (Tibétain) de Qinghai-Xizang, 645  
  
 ombre pluviométrique, 276  
 opération de réservoir, 523  
 optimisation, 432  
   globale, 450  
   locale, 450  
   par essaim de particules, 432, 686  
 Oscillation Nord Atlantique, 74  
 oued, 1103  
   Sebdou, 1271  
 oxygène dissous (OD), 362  
  
 perceptron multi-couches (PMC), 305, 896  
 période de retour, 1216  
 pertes par évapotranspiration, 376  
 plan, 1285  
 plomb-210, 181  
 pluie(s), 917  
   à faibles pas de temps, 1119  
   cumulée, 376  
   extrêmes, 49  
   journalière, 1119  
 pluie-débit, 1103  
   réponse, 843  
 pluviométrie, 863  
 politiques publiques, 1179  
 pollution  
   de l'eau, 702  
   diffuse, 362  
 pompage illégal, 763  
 pré-blanchissement, 611  
 précipitation, 152, 305, 936  
   de mousson estivale, 645  
 prévention, 1016  
 prévisibilité potentielle, 152  
 prévision  
   de crue, 114  
   de débit, 397  
   en bassin non jaugé (PUB), 625  
   hydrologique, 414  
   par réseau de neurones artificiels, 491  
   probabiliste, 523  
   saisonnier dynamique, 152  
   saisonnier, 508  
 principe d'extension, 595  
 processus  
   de Poisson, 917  
   non-linéaire, 793  
   de sédiments, 1253  
   sédimentaire, 1068  
 profondeur équivalente, 338  
 programmation génétique, 563  
 propriétés  
   du sol, 1051  
   spatiales, 1119  
 puissance d'un test, 611  
 puits d'écémage, 352  
  
 qualité de l'eau, 713  
  
 rapport de vraisemblance maximale, 956  
 ré-aération, 362  
 recharge, 1051  
 recuit simulé, 352, 432  
 réflectance hyperspectrale, 808  
 réflectométrie dans le domaine temporel (TDR), 478  
 régime transitoire, 763  
 région aride, 1103  
 régionalisation, 625, 1119  
 régression, 713, 793, 1103  
   régionale, 625  
 regroupement de bassins versants, 625  
 relation hauteur-débit, 595  
 relief, 1088

- réponse
  - des crues, 671
  - en rendement cultural, 686
  - pluie-débit, 843
- réseau(x)
  - à délais temporels, 508
  - Bayésiens, 523
  - de neurones, 141, 414
    - artificiel(s), 352, 397, 713, 563, 793
    - Bayésien, 508
    - simple, 896
  - hydrographique, 230
  - récurifs, 508
- ressources en eau, 152, 1232
  - globales, 247
- restauration lacustre, 671
- rétablissement, 1016
- Rivière
  - Luján, 702
  - Ouse, 414
  - Taquari, 878
  - Tyne (R-U), 992
  - Uruguay, 878
- Rosenbrock, 432
- ruissellement, 1051, 1134
- rupture, 68
- saisonnalité, 1088
- saut, 18, 68
- scénario(s)
  - climatique, 38
  - hydrologiques, 247
- sécheresse(s), 68, 625
- sédimentogramme, 1285
- segmentation de Hubert, 68
- semi-aride tunisien, 1119
- semivariogramme, 1038
- Sénégal, 68
- séparation
  - d'hydrogramme, 491
  - de l'écoulement de base, 491
  - événementielle d'hydrogramme, 843
- série climatique, 18
- shuffled complex evolution, 432
- SIG ArcView, 292
- simplex, 432
- simulation(s), 1119
  - de débit de rivière, 896
- singularité, 68
- site non jaugé, 86
- SMAR, 432
- SOI, 1004
- soulèvement de l'interface, 352
- source thermale, 1298
- source, 192
- Sri Lanka, 1051
- stratification hydrochimique, 732
- stress hydrique, 247
- structure
  - de modèle, 230
  - de propagation sérielle, 114
- sudestadas, 702
- surexploitation, 1146
- SWAT, 450
- système(s)
  - aquifère, 206
  - d'inférence floue, 793
  - hydrogéothermique, 1298
  - neuro-flou, 99
- Taiwan, 114
- taux de sédimentation, 181
- télé-détection, 808
- température de rivière, 74
- temps
  - de transit moyen, 748
  - réel, 318
- tendance(s), 18, 1146
  - à long terme, 74
  - linéaires, 1216
  - temporelles, 1004
- test
  - de Mann-Kendall, 611, 956
  - régional de détection de tendance, 956
  - statistique, 18
- trajectoire pluviogène, 1088
- transposition de données, 86
- Tunisie, 1088, 1162
  - centrale, 1134, 1232, 1298
- turbidigramme, 1068
- types de temps de Lamb, 276
- variabilité
  - climatique, 38
  - interannuelle, 1216
  - spatio-temporelle de la pluie, 230
- variables en zéro-un, 352
- Viêt-nam, 654
- Vulnérabilité, 748
- Zagros, 206
- zone
  - enneigée, 376
  - non saturée, 748
  - semi-aride, 1271
    - tunisienne, 1285
  - tropicale humide, 49

